U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

PB-280 783

Quantum Yields for Reaction of Pollutants in Dilute Aqueous Solution

Environmental Research Lab, Athens, Ga

29 Aug 77

PB 280 783

TECHNICAL REPORT DATA
(Please read Instructions on the reverse before completing)

1. REPORT NO. EPA-600/J-78-004	2. JOURNAL ARTICLE
3. RECIPIENT'S ACCESSION NO.	
4. TITLE AND SUBTITLE Quantum Yields for Reaction of Pollutants in Dilute Aqueous Solution	5. REPORT DATE 6. PERFORMING ORGANIZATION CODE
7. AUTHOR(S) Richard G. Zepp	8. PERFORMING ORGANIZATION REPORT NO.
9. PERFORMING ORGANIZATION NAME AND ADDRESS Environmental Research Laboratory U.S. Environmental Protection Agency College Station Road Athens, GA 30605	10. PROGRAM ELEMENT NO. 1BA609 11. CONTRACT/GRANT NO.
12. SPONSORING AGENCY NAME AND ADDRESS U.S. Environmental Protection Agency-Athens,GA Environmental Research Laboratory College Station Road Athens, GA 30605	13. TYPE OF REPORT AND PERIOD COVERED Journal Article 14. SPONSORING AGENCY CODE EPA/600/01

15. SUPPLEMENTARY NOTES
Published in Environmental Science and Technology 12(3): 327-329,1978
CONTACT: Richard G. Zepp, 8-250-3428 (Athens, GA)

16. ABSTRACT

Procedures are reported for determining quantum yields for direct photolysis of pollutants at low concentrations in water. Quantum yields are used to compute half-lives for direct photolysis of aquatic pollutants under sunlight.

17. KEY WORDS AND DOCUMENT ANALYSIS

a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
Quantum efficiencies		07E
Photolysis		07D
Contaminants		06F
Water		13B
Half-life		18B
Pesticides		68D

REPRODUCED BY
NATIONAL TECHNICAL
INFORMATION SERVICE
U. S. DEPARTMENT OF COMMERCE
SPRINGFIELD, VA. 22161

18. DISTRIBUTION STATEMENT RELEASE TO PUBLIC	19. SECURITY CLASS *(This Report)* UNCLASSIFIED 20. SECURITY CLASS *(This page)* UNCLASSIFIED	21. NO. OF PAGES 5 22. PRICE A02-A01

EPA Form 2220-1 (9-73)

EPA-600/J-78-004
JOURNAL ARTICLE

Quantum Yields for Reaction of Pollutants in Dilute Aqueous Solution

Richard G. Zepp

Environmental Research Laboratory, U.S. Environmental Protection Agency, College Station Road, Athens, Ga. 30605

■ Procedures are reported for determining quantum yields for direct photolysis of pollutants at low concentrations in water. Quantum yields are used to compute half-lives for direct photolysis of aquatic pollutants under sunlight.

In a recent report, Zepp and Cline (*1*) described equations and parameters required to compute direct photolysis rates of pollutants in the aquatic environment. Photolysis rates ($-dC/dt$) can be computed by Equation 1 where k_a is the specific rate of light absorption, ϕ is the quantum yield for reaction, and C is the pollutant concentration.

$$-\frac{dC}{dt} = k_a \phi C \tag{1}$$

Usually, the quantum yield for a reaction is determined under conditions in which all the incident light or a directly measurable fraction is absorbed by the system (*2*). The concentrations of substrate employed in these experiments typically range from 10^{-3} to greater than 0.1 M. The extremely low solubilities of most organic pollutants in water preclude measurement of quantum yields under conditions of complete light absorption. In this communication a technique that can be used to determine quantum yields in very dilute solution is described. An explicit description of the use of quantum yields to compute sunlight photolysis half-lives is also included.

The general rate expression for photolysis in a cell containing volume V of solution and having exposed area A is:

$$\text{Rate} = -\left(\frac{dC}{dt}\right)_\lambda = \phi I_{o\lambda} \left(\frac{A}{V}\right) F_{s\lambda} F_{c\lambda} \tag{2}$$

where $I_{o\lambda}$ is the incident light intensity at wavelength, λ; $F_{s\lambda}$ is the fraction of light absorbed by the system; and $F_{c\lambda}$ is the fraction of absorbed light that is absorbed by the photoreactive substance (*3*). The terms $F_{s\lambda}$ and $F_{c\lambda}$ are expressed by:

$$F_{s\lambda} = 1 - 10^{-(\alpha_\lambda + \epsilon_\lambda C)l} \tag{3}$$

$$F_{c\lambda} = \frac{\epsilon_\lambda C}{\alpha_\lambda + \epsilon_\lambda C} \tag{4}$$

where α_λ is the absorption coefficient of the solvent, ϵ_λ is the molar extinction coefficient of the photoreactive substance, and l is the light pathlength. When light is weakly absorbed by the system ($F_{s\lambda} < 0.1$), $F_{s\lambda}$ very nearly equals $2.303\,(\alpha_\lambda + \epsilon_\lambda C)l$, and Equation 2 simplifies to a first-order rate expression:

$$\text{Rate} = 2.303\, I_{o\lambda} \left(\frac{A}{V}\right) \epsilon_\lambda l \phi C \tag{5}$$

Thus, when the light intensity, extinction coefficient, and pathlength are known, the quantum yield may be computed from the slope of a first-order plot of ln C vs. exposure time:

$$\phi = \frac{-\text{Slope}}{2.303\, I_{o\lambda} (A/V) \epsilon_\lambda l} \tag{6}$$

Procedures that can be used to determine these parameters are outlined below. Monochromatic light isolated by filters (*2*) is usually employed in these experiments.

The intensity term, $I_{o\lambda}(A/V)$, can be directly determined by exposing a solution of an appropriate reference compound, i.e., a chemical actinometer, to light in exactly the same fashion as the pollutant. The same or an identical reaction cell should be used, and the cell should be filled with the same volume. The chemical actinometer should photoreact with a well-defined quantum yield at wavelength λ. When all of the incident light is absorbed by the actinometer, i.e., $F_{s\lambda}$ is unity, the intensity term, $I_{o\lambda}(A/V)$, equals the rate of photoreaction of the actinometer divided by its quantum yield. The potassium ferrioxalate actinometer is reliable and easy to analyze (*4*).

The average pathlength of the photolysis cell can be determined experimentally by the following procedure. If the concentration of the light absorbing substance, C, in a photoreactive system is sufficiently high that $F_{c\lambda}$ (Equation 4) is close to unity, then Equation 2 reduces to:

$$(\text{Rate})_C = \phi I_{o\lambda} \left(\frac{A}{V}\right) (1 - 10^{-\epsilon_\lambda l C}) \tag{7}$$

In the special case in which $\epsilon_\lambda lC$ exceeds 2, essentially all of the light is absorbed, and Equation 7 further simplifies to Equation 8.

$$(\text{Rate})_{\max} = \phi I_{o\lambda} \left(\frac{A}{V}\right) \tag{8}$$

Published 1978 American Chemical Society

The ratio, X, of the rate at concentration C to the maximum rate is then given by:

$$\frac{(\text{Rate})_C}{(\text{Rate})_{\max}} = X = 1 - 10^{-\epsilon_\lambda l C} \tag{9}$$

Thus, a plot of $-\log(1 - X)$ vs. $\epsilon_\lambda C$ gives a straight line with a slope equal to the pathlength, l. Such a plot is shown in Figure 1 for the benzophenone-sensitized *cis-* to *trans-*isomerization of 1,3-pentadiene (*5*) in our photochemical apparatus. In Figure 1, X is the ratio of *trans-*isomer formed at benzophenone concentration C to *trans-*isomer formed at a benzophenone concentration in which all the light was absorbed (0.050 M); ϵ is the extinction coefficient of benzophenone at 313 nm, the wavelength employed in the cell pathlength determination.

If possible, the molar extinction coefficient of the pollutant (ϵ_λ, Equation 3) should be determined in water. This task is sometimes difficult, if not impossible, because the solubility limits of DDT, benzo(a)pyrene, and some other pollutants are lower than 10^{-8} M. As an alternative, we have determined extinction coefficients in mixtures of water and acetonitrile or methanol (*6*), two polar organic solvents whose refractive indices are very close to that of water (*7*). Our choice of organic solvents was dictated mainly by the theoretical prediction that, at least for strong absorption bands of nonpolar compounds, absorption coefficients should be the same for solvents with the same refractive indices (*8*).

In agreement with theory, Schwarz and Wasik (*9*) reported that the absorption spectra of several polycyclic aromatic hydrocarbons are the same in water and ethanol, another organic solvent whose refractive index is nearly the same as that of water. Polar substances are very sensitive to changes in solvent, especially if the substance forms hydrogen bonds with solvent. Because of hydrogen-bonding effects, for example, the spectra of ketones like acetone and of nitrosamines such as *N*-nitrosodimethylamine differ significantly in water and ethanol (*10*). Fortunately, polar substances are usually sufficiently water-soluble that their spectra can be obtained in pure water or in water mixed with only a small amount of organic solvent.

Pure chemicals, not technical grade substances, should be used in the measurement of extinction coefficients. Overestimates of extinction coefficients usually occur when technical substances are employed, because the impurities frequently absorb in the same spectral region as the pure chemical.

Quantum yields may also be determined by comparing the photolysis rates of the pollutant with a reference compound under conditions in which the light is weakly absorbed by both compounds. Such comparisons are conveniently made in the "merry-go-round" photochemical apparatus originally designed by Moses et al. (*11*). The quantum yield for reaction of the pollutant, ϕ, can be calculated:

$$\phi = \frac{(SLOPE)_P}{(SLOPE)_R} \cdot \frac{\epsilon_R \phi_R}{\epsilon_P} \tag{10}$$

where $(SLOPE)_P$ and $(SLOPE)_R$ are the slopes of log plots of concentration vs. exposure time for pollutant and reference compound, respectively; ϵ_R and ϵ_P are molar extinction coefficients of reference compound and pollutant; and ϕ_R is the quantum yield for reaction of the reference compound. Values of $\epsilon_R \phi_R$ at 313 nm for a few pesticides are: DDE in water, 12 (*6, 12*); DDE in hexane, 15 (*6, 12*); DMDE in water, 120 (*6, 12*); DMDE in hexane, 72 (*6, 12*); carbaryl in water (pH 5.5), 2.3 (*13*); parathion in water, 0.39 (*14*); methyl parathion in water, 0.43 (*15*). Unfortunately, most well-established chemical actinometers cannot be analyzed at the low concentrations required for this technique.



Figure 1. Determination of average cell pathlength for quantum yield studies by use of benzophenone-sensitized photoisomerization of *cis*-1,3-pentadiene

Once determined, the quantum yield for reaction can be used to compute the half-life for photolysis of the pollutant under a variety of environmental conditions (*1*). The photolysis half-life is independent of the optical properties of a water body near its surface (*1*). Estimates of the effects of light attenuation in water bodies appear elsewhere (*1*). Near surface half-lives can be computed by Equation 11 where Z_λ is underwater solar irradiance (*1*) at a wavelength interval centered at λ, and $\bar{\epsilon}_\lambda$ is the average extinction coefficient at this wavelength interval. The value of j is 6.02×10^{20} photon mole^{-1}.

$$t_{1/2} = \frac{0.693}{k_a \phi} = \frac{0.693\, j}{2.303\, \phi \sum \bar{\epsilon}_\lambda Z_\lambda} \tag{11}$$

A sample computation of the half-life of the pesticide carbaryl is presented in Table I. In computing sunlight half-lives, we have generally assumed that the quantum yield is wavelength independent in the region of sunlight absorption (*1*). This assumption is valid for most complex molecules in solution because photoreaction from second or higher electronic states usually cannot compete with the rapid radiationless decay of excited molecules to their first excited state. If photoreaction often occurred from higher excited states, excitation spectroscopy would not have general applicability (*16*). Exceptions to the general rule include certain coordination compounds. For example, the photolysis of iron (II) cyanide complexes is clearly wavelength dependent (*3*). For such cases, sunlight half-lives equal 0.693 $(\Sigma \phi_\lambda \epsilon_\lambda Z_\lambda)^{-1}$ where ϕ_λ is the average quantum yield in the wavelength interval centered at λ.

Finally, it should be emphasized that direct photolysis is but one of several potential mechanisms for photochemical transformation of pollutants. Natural substances in the aquatic environment are known to sometimes accelerate photolysis by mechanisms that are presently not defined (*6, 17*). Nonetheless, several studies (*6, 13, 15*) indicate that photolysis half-lives determined in natural waters are often nearly the same as half-lives computed by the procedures described above.

Literature Cited

(1) Zepp, R. G., Cline, D. M., *Environ. Sci. Technol.*, **11**, 359 (1977).
(2) Calvert, J. G., Pitts, J. N., "Photochemistry", pp 686–798, Wiley, New York, N.Y., 1966.
(3) Balzani, V., Carassiti, V., "Photochemistry of Coordination Compounds", pp 6–14, 146–51, Academic Press, New York, N.Y., 1970.
(4) Hatchard, C. G., Parker, C. A., *Proc. Roy. Soc.* (*London*), **A235**, 518 (1956).

Table I. Sample Calculations of Direct Photolysis Half-Life of Carbaryl in Water Midday, Midsummer, Latitude 40°N

Wavelength, nm	ϵ_λ	$Z_\lambda \times 10^{-14}$	$\epsilon_\lambda Z_\lambda \times 10^{-15}$
297.5	1480	0.00716	1.06
300	918	0.0240	2.20
302.5	741	0.0723	5.36
305	532	0.181	9.63
307.5	427	0.305	13.0
310	356	0.495	17.6
312.5	388	0.717	27.8
315	261	0.933	24.3
317.5	235	1.15	27.0
320	101	1.35	13.6
323	45	2.52	11.3
330	11	8.46	9.31
			$\sum \bar{\epsilon}_\lambda Z_\lambda = 162$

$$k_a = \frac{2.303}{6.02 \times 10^{20}} \times 1.62 \times 10^{17} = 6.20 \times 10^{-4}\ s^{-1}$$

$$\phi_{313} = 0.0060$$

$$t_{1/2} = \frac{0.693}{(0.0060 \times 6.20 \times 10^{-4})} = 1.86 \times 10^5\ s = 51.7\ h$$

(5) Lamola, A. A., Hammond, G. S., *J. Chem. Phys.*, **41**, 2129 (1965).
(6) Zepp, R. G., Wolfe, N. L., Gordon, J. A., Fincher, R. C., *J. Agric. Food Chem.*, **26**, 727 (1976).
(7) Riddick, J. A., Bunger, W. B., "Organic Solvents, Physical Properties and Methods of Purification", Wiley-Interscience, New York, N.Y., 1970.
(8) Strickler, S. J., Berg, R. A., *J. Chem. Phys.*, **37**, 814 (1962).
(9) Schwarz, R. P., Wasik, S. P., *Anal. Chem.*, **48**, 524 (1976).
(10) Jaffé, H. H., Orchin, M., "Theory and Applications of Ultraviolet Spectroscopy", pp 186–8, Wiley, New York, N.Y., 1962.
(11) Moses, F. G., Liu, R.S.H., Monroe, B. M., *Mol. Photochem.*, **1**, 245 (1969).
(12) Zepp, R. G., Wolfe, N. L., Azarraga, L. V., Cox, R. H., Pape, C. W., *Arch. Environ. Contam. Toxicol.*, **6**, 305 (1977).
(13) Wolfe, N. L., Zepp, R. G., Baughman, G. L., Fincher, R. C., Gordon, J. A., "Chemical and Photochemical Transformation of Selected Pesticides in Aquatic Systems", EPA Report No. EPA-600/3-76-067, Sept. 1976.
(14) Zepp, R. G., unpublished results.
(15) Smith, J. H., Mabey, W. R., Bohonos, N., Holt, B. R., Lee, S. S., Chou, T.-W., Bomberger, D. C., Mill, T., "Environmental Pathways of Selected Chemicals in Freshwater Systems", EPA Report No. EPA-600/7-77-113, Oct. 1977.
(16) Parker, C. A., in "Advances in Photochemistry", W. A. Noyes, G. S. Hammond, and J. N. Pitts, Vol 11, pp 308–9, Interscience, London, England, 1964.
(17) Ross, R., Crosby, D. G., *Chemosphere*, **4**, 277 (1975).

Received for review May 6, 1977. Accepted August 29, 1977.